



Publications with Simi BioCell

Nehme R, Grote P, Tomasi T, Löser S, Holzkamp H, Schnabel R, Conrath B. (2010)
Transcriptional upregulation of both egl-1 BH3-only and ced-3 caspase is required for the death of the male-specific CEM neurons.
Cell Death Differ. Feb 12.

Cabello J, Neukomm LJ, Günesdogan U, Burkart K, Charette SJ, Lochnit G, Hengartner MO, Schnabel R. (2010)
The Wnt pathway controls cell death engulfment, spindle orientation, and migration through CED-10/Rac.
PLoS Biol. Feb 2;8(2):e1000297

Morris SA, Teo RT, Li H, Robson P, Glover DM, Zernicka-Goetz M. (2010)
Origin and formation of the first two distinct cell types of the inner cell mass in the mouse embryo.
Proc Natl Acad Sci U S A. Apr 6;107(14):6364-9.

Jedrusik A, Bruce AW, Tan MH, Leong DE, Skamagki M, Yao M, Zernicka-Goetz M. (2010)
Maternally and zygotically provided Cdx2 have novel and critical roles for early development of the mouse embryo.
Dev Biol. Apr 27. [Epub ahead of print]

Bertrand V, Hobert O. (2009)
Linking asymmetric cell division to the terminal differentiation program of postmitotic neurons in *C. elegans*.
Dev Cell. Apr;16(4):563-75.

Lahl V, Schulze J, Schierenberg E. (2009)
Differences in embryonic pattern formation between *Caenorhabditis elegans* and its close parthenogenetic relative *Diploscapter coronatus*.
Int J Dev Biol.;53(4):507-15.

Sarin S, Antonio C, Tursun B, Hobert O. (2009)
The *C. elegans* Tailless/TLX transcription factor nhr-67 controls neuronal identity and left/right asymmetric fate diversification.
Development. Sep;136(17):2933-44. Epub 2009 Jul 29.

Bischoff M, Cseresnyés Z. (2009)
Cell rearrangements, cell divisions and cell death in a migrating epithelial sheet in the abdomen of *Drosophila*.
Development. Jul;136(14):2403-11.

Langenhan T, Prömel S, Mestek L, Esmaeili B, Waller-Evans H, Hennig C, Kohara Y, Avery L, Vakonakis I, Schnabel R, Russ AP. (2009)
Latrophilin signaling links anterior-posterior tissue polarity and oriented cell divisions in the *C. elegans* embryo.
Dev Cell. Oct;17(4):494-504.

Asencio C, Navas P, Cabello J, Schnabel R, Cypser JR, Johnson TE and Rodríguez-Aguilera C (2009)
Coenzyme Q supports distinct developmental processes in *Caenorhabditis elegans*
Mechanisms of ageing and Development, 130, 145-153

Meyvis Y, Houthoofd W, Visser A, Borgonie G, Gevaert K, Vercruyse J, Claerebout E, Geldhof P (2009)
Analysis of the translationally controlled tumour protein in the nematode *Ostertagia ostertagi* and *Caenorhabditis elegans* suggests a pivotal role in egg production
Int J Parasitol, 39(11):1205-13.

Moser SC, von Elsner S, Büssing I, Alpi A, Schnabel R, Gartner A (2009)
Functional Dissection of *Caenorhabditis elegans* CLK-2/TEL2 Cell Cycle Defects during Embryogenesis and Germline Development
Plos Genetics, 5, e1000451

Willems M, Egger B, Wolff C, Mouton S, Houthoofd W, Fonderie P, Couvreur M, Artois T, Borgonie G. (2009)

Embryonic origins of hull cells in the flatworm *Macrostomum lignano* through cell lineage analysis: developmental and phylogenetic implications.

Dev Genes Evol. 2009 Oct 16 [Epub ahead of print].

Bertrand V, Hobert O. (2009)

Linking asymmetric cell division to the terminal differentiation program of postmitotic neurons in *C. elegans*.

Dev Cell;16(4):563-75.

Schulze J, Schierenberg E. (2009)

Embryogenesis of *Romanomermis culicivorax*: an alternative way to construct a nematode.

Dev Biol. 334:10-21.

Bischoff M, Parfitt DE, Zernicka-Goetz M (2008)

Formation of the embryonic-abembryonic axis of the mouse blastocyst: relationships between orientation of early cleavage divisions and pattern of symmetric/asymmetric divisions.

Development. 135, 953-62. Epub 2008 Jan 30.

Houthoofd W, Willems M, Jacobsen K, Coomans A, Borgonie G (2008)

The embryonic cell lineage of the nematode *Rhabditophanes* sp.

Int J Dev Biol, 5, 963-7.

Jedrusic A, Parfitt DE, Guo G, Skamagki M, Grabarek JB, Johnson MH, Robson P, Zernicka-Goetz M (2008)

Role of *Cdx2* and cell polarity in cell allocation and specification of trophoderm and inner cell mass in the mouse embryo

Genes Dev, 22, 2692-706.

Ray P, Schnabel R and Okkema PG (2008)

Behavioral and synaptic defects in *C. elegans* lacking the NK-2 homeobox gene *ceh-28*

Developmental Neurobiology, 68, 421- 433

R. Ravin, D. Hoepfner, D. Munno, L. Carmel, J. Sullivan, D. Levitt, J. Miller, C. Athaide, D. Panchision, R. McKay (2008)

Potency and Fate Specification in CNS Stem Cell Populations In Vitro

Cell Stem Cell, 3, Issue 6, Pages 670-680

Schulze J, Schierenberg E (2008)

Cellular pattern formation, establishment of polarity and segregation of colored cytoplasm in embryos of the nematode *Romanomermis culicivorax*

Dev Biol, 315, 426-36. Epub 2008 Jan 8.

Smit RB, Schnabel R, Gaudet J (2008)

The HLH-6 transcription factor regulates *C.elegans* pharyngeal gland development and function

PLoS Genetics, 4, e1000222. Epub 2008 Oct 17.

Stach T, Winter J, Bouquet JM, Chourout D, Schnabel R (2008)

Embryology of a planktonic tunicate reveals traces of sessility

Proc Natl Acad Sci U S A. 105, 7229-34. Epub 2008 May 19. Erratum in: Proc Natl Acad Sci U S A.2008 105,10268.

von Hilchen CM, Beckervordersandforth RM, Rickert C, Technau GM, Altenhein B (2008)

Identity, origin, and migration of peripheral glial cells in the *Drosophila* Embryo

Mech Dev.125, 337-52. Epub 2007 Nov 1.

Hunt-Newbury R, Viveiros R, Johnsen R, Mah A, Anastas D, Fang L, Halfnight E, Lee D, Lin J, Lorch A, McKay S, Okada HM, Pan J, Schulz AK, Tu D, Wong K, Zhao Z, Alexeyenko A, Burglin T, Sonnhammer E, Schnabel R, Jones SJ, Marra MA, Baillie DL, Moerman DG (2007)

High-throughput in vivo analysis of gene expression in *Caenorhabditis elegans*.

PLoS Biology, 5, e237

Tiraihi A, Tiraihi (2007)

Early onset of regionalization in EMS lineage of *C. elegans* embryo: a quantitative study.

Biosystems, 90, 676-86. Epub 2007 Mar 3

Asencio C, Rodríguez-Aguilera JC, Vázquez R, Baylis H, Cabello J, Schnabel R, Gavilán A, Navas P (2006)
Differential expression pattern of *coq-8* gene during development in *Caenorhabditis elegans*
Gene Expr Patterns, 6, 433-9.

Ralf Schnabel, Marcus Bischoff, Arend Hintze, Anja-Kristina Schulz, Hans Meinhardt and Harald Hutter (2006)
Global cell sorting in the *C. elegans* embryo defines a new mechanism for pattern formation.
Developmental Biology, 294, 418-431

Bischoff M and Schnabel R (2006)
A posterior centre establishes and maintains polarity of the *Caenorhabditis elegans* embryo by a Wnt-dependent relay mechanism.
PLoS Biol, 4,e396.

Bischoff M and Schnabel R (2006)
Global cell sorting is mediated by local cell-cell interactions in the *C. elegans* embryo.
Developmental Biology, 294, 432-444

Hejnol A, Schnabel R, Scholtz G (2006)
A 4D-microscopic analysis of the germ band in the isopod crustacean *Porcellio scaber* (Malacostraca, Peracarida)—developmental and phylogenetic implications.
Development Genes and Evolution, 216, 755-767

Wolff C, Scholtz G (2006)
Cell lineage analysis of the mandibular segment of the amphipod *Orchestia cavimana* reveals that the crustacean paragnaths are sternal outgrowths and not limbs.
Front Zool, 4,19

Houthoofd W, Willems M, Vangestel S, Mertens C, Bert W, Borgonie G (2006)
Different roads to form the same gut in nematodes.
Evol Dev, 8, 362-9.

Gavilán A, Asencio C, Cabello J, Rodríguez-Aguilera JC, Schnabel R, Navas P (2005).
C. elegans knockouts in ubiquinone biosynthesis genes result in different phenotypes during larval development.
Biofactors. 2005;25, 21-9.

Kinchen JM, Cabello J, Klingele D, Feichtinger R, Schnabel H, Schnabel R and Hengartner MO (2005)
Two pathways converge at CED-10 to mediate actin rearrangement and corpse removal in *C. elegans*
Nature, 434, 93-99

Hejnol A & Schnabel R (2005)
The eutardigrade *Thulinia stephaniae* has an indeterminate development and the potential to regulate early blastomere ablations.
Development, 132, 1349-1361

Gavilán A, Asencio C, Cabello J, Rodríguez-Aguilera JC, Schnabel R, Navas P (2005)
C. elegans knockouts in ubiquinone biosynthesis genes result in different phenotypes during larval development
Biofactors, 25, 21-9

Braun V, Azevedo RB, Gumbel M, Agapow PM, Leroi AM, Meinzer HP (2003).
ALES: cell lineage analysis and mapping of developmental events
Bioinformatics, 19, 851-8.

Ian A. Hope, Andrew Mounsey, Petra Bauer and Sobia Aslam (2003)
The forkhead gene family of *Caenorhabditis elegans*
Gene 304, 43-55

Houthoofd W, Jacobsen K, Mertens C, Vangestel S, Coomans A, Borgonie G (2003)
Embryonic cell lineage of the marine nematode *Pellioditis marina*.
Dev Biol, 258, 57-69

Urbach R, Schnabel R and Technau GM (2003)

The pattern of neuroblast formation, mitotic domains and proneural gene expression during early brain development in *Drosophila*.

Development, 130, 3589-3606

Karabinos A, Büssing, I, Schulz E, Wan, J, Weber K and Schnabel R. (2003)

Functional analysis of the single calmodulin gene in the nematode *C. elegans* by RNA interference and 4-D microscopy

European. J. Cell Biology, 82, 557-563 (82 citations)

Clucas C, Cabello J, Büssing I, Schnabel R, Johnstone IL (2002)

Oncogenic potential of a *C.elegans cdc25* gene is demonstrated by a gain-of- function allele

EMBO J. 21, 665–674.

Ginzburg VE, Roy PJ, Culotti JG (2002)

Semaphorin 1a and semaphorin 1b are required for correct epidermal cell positioning and adhesion during morphogenesis in *C.elegans*.

Development, 129, 2065-78

Hoepfner DJ, Hengartner MO, Schnabel R (2001)

Engulfment genes cooperate with *ced-3* to promote cell death in *Caenorhabditis elegans*.

Nature, 412, 202-206

Gaëtan Borgonie, Kim Jacobsen & August Coomans (2000)

Embryonic lineage evolution in nematodes

Nematology, 2, 65-69

Dantoni JC, Quintin S, Lakatos L, Labouesse M, Tora L (2000)

TBP-like factor is required for embryonic RNA polymerase II transcription in *C. elegans*.

Mol Cell, 6, 715-22.

Brunschwig K, Wittmann C, Schnabel R, Bürglin TR, Tobler H, Müller F (1999)

Anterior organization of the *Caenorhabditis elegans* embryo by the labial-like Hox gene *ceh-13*

Development, 126,1537-46

Schnabel R (Zur Verfügung gestellt)

Biology in pictures: a question of fate (1999)

Curr. Biol. 12, R543.

Schnabel R (1999)

Microscopy in *C. elegans*

A Practical Approach (I.A. Hope, ed.) 119-141

Dolinski C, Borgonie G, Schnabel R, Baldwin JG (1998)

Buccal capsule development as a consideration for phylogenetic analysis of Rhabditida (Nemata).

Dev Genes Evol, 208, 495-503.

Horner MA, Quintin S, Domeier ME, Kimble J, Labouesse M, Mango SE (1998)

pha-4, an HNF-3 homolog, specifies pharyngeal organ identity in *Caenorhabditis elegans*.

Genes Dev, 12,1947-52

Schnabel R & Schnabel H (1997)

Hox genes misled by local environments

Nature, 385, 588-589

Kaletta T, Schnabel H & Schnabel R (1997)

Binary specification of the embryonic lineage in *Caenorhabditis elegans*

Nature, 390, 294-298

Schnabel R, Hutter H, Moerman D, Schnabel H (1997)

Assessing normal embryogenesis in *Caenorhabditis elegans* using a 4D microscope: variability of development and regional specification.

Dev Biol,184, 234-65.